



## After Port-a-Cath Removal

1. Keep the dressing dry for \_\_\_\_\_ hours. You may remove it after \_\_\_\_\_ hours.
2. After removing the dressing, you may shower. Wash the incision with soap and water, then pat dry.
3. If you see drainage at the site, cover the site with a dressing as needed. Remove and change the dressing each day.
4. If “Steri-strips” are used, leave them in place until they fall off (about 7 to 10 days).
5. If Dermabond is used, there will be no dressing. Dermabond will stay in place for 5 to 10 days; then fall off on its own.
6. Limit activity involving your arms for the next 72 hours. Do no strenuous exercise or activity for 1 week.
7. If you have pain, use Tylenol, Motrin, or Advil, if no other pain medication was prescribed.
8. If you have any of the following signs of infection, contact your medical staff fellow via the NIH page operator at **301-496-1211**:
  - fever
  - chills
  - increased redness and tenderness at the site
  - excessive drainage from the site

Other Instructions:

**Contacts** (Reach the staff members below through the NIH page operator at **301-496-1211**.)

Your Doctor:


This information is prepared specifically for persons taking part in clinical research at the National Institutes of Health Clinical Center and may not apply to patients elsewhere. If you have questions about the information presented here, talk to a member of your health-care team.

Products/resources named serve as examples and do not imply endorsement by NIH. The fact that a certain product/resource is not named does not imply that such product/resource is unsatisfactory.

National Institutes of Health Clinical Center  
Bethesda, MD 20892

Questions about the Clinical Center?  
<http://www.cc.nih.gov/comments.shtml>  
12/07

